

▼ This medicine is subject to additional monitoring.

If you get any side effects, talk to your doctor, pharmacist or nurse.

You can also report side effects directly. Report forms can be found at

**[www.medicinesauthority.gov.mt/adrportal](http://www.medicinesauthority.gov.mt/adrportal)** and sent by email to [postlicensing.medicinesauthority@gov.mt](mailto:postlicensing.medicinesauthority@gov.mt) or posted to ADR Reporting,

The Medicines Authority Post-licensing Directorate, 203, Level 3,

Rue D'Argens, Gżira GŻR 1368, MALTA. Side effects should also be reported to Roche Products Ltd. Please contact Roche Drug Safety Centre by emailing [welwyn.uk\\_dsc@roche.com](mailto:welwyn.uk_dsc@roche.com) or calling +44 (0)1707 367554.



### Emergency Contact Phone Numbers

Office Hours

--

Out of Hours

--

**Erivedge<sup>®</sup>▼ (vismodegib)**

*Patient reminder card*

Marketing authorisation holder: Roche Registration Ltd, 6 Falcon Way, Shire Park, Welwyn Garden City, AL7 1TW, United Kingdom. Erivedge is a registered trademark.

RXUKERIV00077c

Date of preparation: November 2013

This is produced by Roche Products Ltd.

**Erivedge may harm or cause the death of an unborn child.**

**FEMALE PATIENTS** who are able to become pregnant must:

- always use **2** forms of recommended contraception during treatment and for 24 months after your last dose (please refer to the “Erivedge® Pregnancy Prevention Programme: Important information for men and women taking Erivedge about pregnancy prevention and contraception” for more information)
- have monthly pregnancy tests during treatment to monitor for pregnancy

**MALE PATIENTS** must:

- always use a condom (with spermicide, if available) when having sex with female partners during treatment and for 2 months from last dose
- not donate semen during treatment and for 2 months after the final dose

You must talk to your healthcare professional immediately if you or your sexual partner misses a period, suspect a pregnancy, or are pregnant.

- **FEMALE PATIENTS:** talk to your doctor if you suspect a pregnancy while taking Erivedge and for 24 months after your final dose.
- **MALE PATIENTS:** talk to your doctor if your female partner suspects that she is pregnant while you are taking Erivedge and for 2 months after your final dose.

**ALL PATIENTS** must:

- never give this medicine to anyone else
- not donate blood during treatment and for 24 months after the final dose
- return all unused capsules at the end of treatment