HMA WGEO – Rapid Alert Form Counterfeit or illegal product found in the illegal supply chain

Reference:				
Date:	Time:		Initials:	
Please complete sections 1 to 5 providing as much information as possible.				
1. REPORTING PERSO	ON	ph.		
Name:		Position:		
Organisation:				
Address:			4	
Telephone No:			Ext:	
e-mail address:				
2. PRODUCT DETAILS				
Product name: Equipos	e®			
Manufacturer: Meditech Human Pharmaceuticals, Lörachstr. 278, Ingoldheim				
Supplier: Ilke Altintas				
Ozgürlük mh. Mrt				
Antalya, Türkei				
Legal status: Banned	- A - A - A - A - A - A - A - A - A - A	Unlicense	ed Stolen	
Dosage form: Injection				
Strength: 250 mg				
Batch / lot no: C455				
If yes to the above, advise batch destination country:				
Expiry date: 02/21				
Language of packaging	: English			
Date of discovery: 15th Feb. 2019				
Details of discovery: R customs (import by unauth			e received the information via	
Meditech Human Pharmaceutical is unknown;				
Ingoldheim is not located in Germany.				
Analysed: YES ☐ NO ☒				
If yes, result of analysis	1			

	10.			
Internet: YES 🛛 / NO 🗌	28			
Internet:	Non internet, advise full details:			
URL:				
Website address:				
https://www.meditechpharmaceutical.net/				
Other details:				
Currency of payment:				
Has product reached patients/consumers?				
4. RISK TO PUBLIC HEALTH				
Adverse reactions: YES / NO				
If yes, please advise details:				
Medical assessment details:				
5. NEED FOR PUBLICITY				
Are you making a public statement? YES ☐ / NO ☒				
Are you issuing a press release? YES□ / NO ⊠				
Are you recalling product? YES ☐ / NO ☒				
If yes to any of the above, when do you intend to take action?				
6. DISSEMINATION				
Are you content for this Rapid Alert to be shared outside WGEO membership?				
YES ☐ / NO ☐ (please see below)				
If yes, please specify which of the below you are content for this to be shared				
with (you may tick more than 1 box)				
Law Enforcement Industry Security Trade Associations				
Traders Other Please specify				
7. PHOTOGRAPH				
See attachment				

3. DISTRUBUTION METHOD















DECANAL

Nandrolone Decanoate is an anabolic steroid. Nandrolone decanoate is used in the treatment of osteoporosis in postmenopausal women (though now not recommended) at a dose of 50 mg every three weeks. It is also used for some aplastic anemias. The positive effects of the drug include muscle growth, appetite stimulation and increased red blood cell production and bone density. Clinical studies have shown it to be effective in treating anemia, osteoporosis and some forms of neoplasia including breast cancer, and also acts as a progestin-based contracentive. progestin-based contraceptive.

Nandrolone is not broken down into DHT, the deleterious effects common to most anabolic steroids on the scalp, skin, and prostate are lessened to a degree; but is rather broken down to the much weaker androgen dihydronandrolone. The drug's fiver toxicity. Estrogenic effects resulting from reaction with aromatase are also recided due to lessened enzyme interaction but effects such as gynaecomastic and reduced libido still occur in larger doses because of other mechanisms. Other side-effects of abuse can include erectile dysfunction and cardiovascular damage, as well as several ailments resulting from the drug's effect of lowering levels of luteinizing hormone through negative feedback. Erectile dysfunction is attributed to the weaker action of dihydronandrolone in the penis since dihydrotestosterone is a known sexual modulator.

ANABOLIC/ANDROGENIC RATE
Anabolic/Androgenic rate of Nandrolone Decanoate is 125:37

DOSAGE For men: 200-800mg/week. For women: 50-100mg/week.

HALF-LIFE

Half-Life of Nandrolone Decandate is 15 days.

DETECTION TIME

Nandrolone Decanoate detection time is 18 months.































Keep out of the reach and

sight of childiren. Do not use after the expiry date.

DOSAGE

For dosage, administration & detailed direction for use, see package insert.

FOR INTRAMUSCULAR USE

To be sold by retail on the prescription of a Registered Medical Practitioner only

STORAGE

Do not refrigerate or freeze. Do not store above 30°C Protect from light.

WARNING

Keep out of the reach and sight of childiren. Do not use after the expiry date.

DOSAGE

For dosage, administration & detailed direction for use, see package insert.

FOR INTRAMUSCULAR USE

To be sold by retail on the prescription of a Registered Medical Practitioner only

















